



PubMed	Nucleotide	Protein	Genome	Structure	PMC	Taxonomy	OMIM	Books
Search PubMed	<input type="text" value="for"/>						Go	Clear
Limits		Preview/Index		History		Clipboard		Details

Display	Abstract	<input type="text" value="20"/>	Show:	20	Sort	<input type="text" value=""/>	Send to	Text	<input type="text" value=""/>
---------	----------	---------------------------------	-------	----	------	-------------------------------	---------	------	-------------------------------

1: Science 1995 Oct 20;270(5235):467-70

[Related Articles, Links](#)

Comment in:

- Science. 1995 Oct 20;270(5235):368-9, 371.

Quantitative monitoring of gene expression patterns with a complementary DNA microarray.

Schena M, Shalon D, Davis RW, Brown PO.

Department of Biochemistry, Beckman Center, Stanford University Medical Center, CA 94305, USA.

A high-capacity system was developed to monitor the expression of many genes in parallel. Microarrays prepared by high-speed robotic printing of complementary DNAs on glass were used for quantitative expression measurements of the corresponding genes. Because of the small format and high density of the arrays, hybridization volumes of 2 microliters could be used that enabled detection of rare transcripts in probe mixtures derived from 2 micrograms of total cellular messenger RNA. Differential expression measurements of 45 Arabidopsis genes were made by means of simultaneous, two-color fluorescence hybridization.

PMID: 7569999 [PubMed - indexed for MEDLINE]

Display	Abstract	<input type="text" value="20"/>	Show:	20	Sort	<input type="text" value=""/>	Send to	Text	<input type="text" value=""/>
---------	----------	---------------------------------	-------	----	------	-------------------------------	---------	------	-------------------------------

[Write to the Help Desk](#)
[NCBI](#) | [NLM](#) | [NIH](#)
[Department of Health & Human Services](#)
[Freedom of Information Act](#) | [Disclaimer](#)